arXiv:1610.03315 KA–TP–34–2016 (v2)

Lorentz-violating neutral-pion decays in isotropic modified Maxwell theory

F.R. Klinkhamer*

Institute for Theoretical Physics, Karlsruhe Institute of Technology (KIT), 76128 Karlsruhe, Germany

arXiv:1610.03315v2 [hep-ph] 27 Oct 2016

Abstract

We consider an extension of the Standard Model with isotropic nonbirefringent Lorentz violation in the photon sector and restrict the discussion to the case of a "fast" photon with a phase velocity larger than the maximum attainable velocity of the fermions. With our conventions, this case corresponds to a negative Lorentz-violating parameter κ in the action. The decay rate of a neutral pion into two photons is calculated as a function of the 3-momentum of the initial pion and the negative Lorentz-violating parameter κ of the final photons.

PACS numbers: 11.30.Cp, 12.20.-m, 13.85.Tp, 98.70.Sa

* frans.klinkhamer@kit.edu

In this note, we calculate the decay rate of a particular Lorentz-violating process, neutral-pion decay into two photons, which can be tested by modeling and observing extensive air showers in the upper Earth atmosphere.

Consider the Standard Model action density augmented by a single Lorentz-violating CPT-even dimension-4 photonic term [1–4] with a dimensionless coupling constant κ. The Lorentz-violating (LV) parameter κ is taken to be negative and to have a very small magnitude (a possible underlying theory has been given in Ref. [5]). A photon of 3-momentum $\vec{k}$ then has a modified dispersion relation,

$$\omega(\vec{k})^2 = \frac{1-\kappa}{1+\kappa}\, c^2\, |\vec{k}\,|^2\,, \tag{1}$$

and modified polarization vectors,

$$\vec{e}^{\,(1)}(\vec{k}\,) \;=\; \sqrt{\frac{1}{1+\kappa}}\, \mathcal{R}_{\widehat{k}} \cdot \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}\,, \tag{2a}$$

$$\vec{e}^{\,(2)}(\vec{k}\,) \;=\; \sqrt{\frac{1}{1+\kappa}}\, \mathcal{R}_{\widehat{k}} \cdot \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}\,, \tag{2b}$$

with the 3×3 rotation matrix $\mathcal{R}_{\widehat{k}}$ which transforms the unit column vector $(1,0,0)^{\mathrm{T}}$ to the unit column vector $\widehat{k} \equiv \vec{k}/k \equiv \vec{k}/|\vec{k}\,|$. See Ref. [4] for further details. Also note that we employ the Minkowski metric $g_{\mu\nu}(x) = \eta_{\mu\nu} \equiv [\mathrm{diag}(+1,\,-1,\,-1,\,-1)]_{\mu\nu}$ and that, from now on, we use natural units with $\hbar = c = 1$. Incidentally, the velocity c corresponds to the maximum attainable velocity of the quarks and leptons.

In this theory, we study the decay of a neutral pion (π^0) of mass M into two nonstandard photons ($\widetilde{\gamma}$), with energies and 3-momenta denoted as follows:

$$\pi^0(E,\,\vec{q}\,) \to \widetilde{\gamma}(\omega,\,\vec{k}\,) + \widetilde{\gamma}(\omega',\,\vec{k}\,')\,, \tag{3}$$

for on-shell energies $E(\vec{q}\,) = \sqrt{|\vec{q}\,|^2 + M^2}$ and $\omega(\vec{k}\,) \geq 0$ from (1). This LV decay process has already been discussed qualitatively in Ref. [6]. A later heuristic discussion in terms of effective-mass-squares [7] has been given in App. A of Ref. [8].

The general expression for the neutral-pion decay parameter γ [7] is as follows (see, e.g., Sec. 3.6 of Ref. [9]):

$$\begin{aligned} \gamma(\vec{q}\,) \;&\equiv\; 2\,\omega(\vec{q}\,)\,\Gamma(\vec{q}\,) \\ &=\; \frac{1}{2}\,\frac{1}{(2\pi)^2} \int \frac{d^3k}{2\,\omega(\vec{k}\,)} \int \frac{d^3k'}{2\,\omega(\vec{k}\,')}\, \delta^4(q-k-k')\, |\mathcal{M}|^2\,, \end{aligned} \tag{4}$$

with the symmetry factor 1/2 and the matrix element $\mathcal{M}$. In a Lorentz-invariant theory, $|\mathcal{M}|^2$ is a scalar and the right-hand side of (4) is manifestly Lorentz invariant, so that γ becomes independent of $\vec{q}$ and is called the decay constant.

Following the discussion of Secs. 4.5 and 7.1 in Ref. [9], we now take the effective pion-photon-photon interaction to be given by

$$\mathcal{L}_{\text{eff}} = \alpha\, C\, \phi\, \epsilon^{\alpha\beta\gamma\delta}\, F_{\alpha\beta}\, F_{\gamma\delta}\,, \tag{5}$$

with the neutral-pion pseudoscalar field ϕ, the Maxwell field strength $F_{\mu\nu} \equiv \partial_\mu A_\nu - \partial_\nu A_\mu$, and a coupling constant C of mass dimension -1. A straightforward but tedious calculation of the LV decay parameter (4) then gives

$$\gamma(\vec{q}) \;=\; \frac{1}{2}\,\frac{1}{8\pi}\,\left(8\,\alpha\, C\right)^2\,\frac{1}{2}\,M^4\; g\left(\frac{q}{M},\,\kappa\right)\,, \tag{6}$$

in terms of a dimensionless function g of the 3-momentum modulus $q \equiv |\vec{q}|$ in units M and the LV parameter κ. The calculated function $g(q/M,\,\kappa)$ has the following properties:

$$g(q/M,\,0) \;=\; 1\,, \tag{7a}$$

$$g(0,\,\kappa) \;=\; \frac{\sqrt{1-\kappa^2}}{(1-\kappa)^3} = 1 + 3\,\kappa + \mathrm{O}(\kappa^2)\,, \tag{7b}$$

$$g(q/M,\,\kappa) \;=\; 0 \;\text{ for }\; \sqrt{q^2+M^2} \geq E^{\,(\text{cutoff})}\,, \tag{7c}$$

with cutoff energy

$$E^{\,(\text{cutoff})} = \sqrt{\frac{1-\kappa}{-2\,\kappa}}\; M \sim \frac{M}{\sqrt{-2\,\kappa}}\,. \tag{8}$$

The last expression for the cutoff energy in (8) agrees with the kinematic result of Ref. [6].

The exact formula for $g(q/M,\,\kappa)$ is, at first, rather cumbersome but can be brought to a manageable form,

$$g\left(\frac{q}{M},\,\kappa\right) \;=\; \begin{cases} \dfrac{\sqrt{1-\kappa^2}}{(1-\kappa)^3}\,\Big[1-(q/q_c)^2\Big]^2 & \text{for }\; q < q_c\,, \\ 0 & \text{for }\; q \geq q_c\,, \end{cases} \tag{9a}$$

$$q_c \;=\; \sqrt{\frac{1+\kappa}{-2\,\kappa}}\; M \sim \frac{M}{\sqrt{-2\,\kappa}}\,. \tag{9b}$$

Figure 1 gives a plot of the function $g(q/M,\,\kappa)$ for a relatively large value of the Lorentz-violating parameter κ.

The final result (9) for the Lorentz-violating neutral-pion decay parameter (6) can be used in numerical simulations of extensive air showers, as discussed in Ref. [8].

ACKNOWLEDGMENTS

It is a pleasure to thank M. Risse for many discussions over the last years. This work was supported in part by the German Research Foundation (DFG) under Grant No. KL 1103/4-1.



FIG. 1. Function $g(q/M,\, \kappa)$ for the neutral-pion decay parameter (6). The numerical value of the Lorentz-violating parameter κ is taken to be $\kappa = -0.5 \times 10^{-4}$.

[1] S. Chadha and H.B. Nielsen, "Lorentz invariance as a low-energy phenomenon," Nucl. Phys. B **217**, 125 (1983).

[2] V.A. Kostelecký and M. Mewes, "Signals for Lorentz violation in electrodynamics," Phys. Rev. D **66**, 056005 (2002), arXiv:hep-ph/0205211.

[3] F.R. Klinkhamer and M. Schreck, "New two-sided bound on the isotropic Lorentz-violating parameter of modified Maxwell theory," Phys. Rev. D **78**, 085026 (2008), arXiv:0809.3217.

[4] F.R. Klinkhamer and M. Schreck, "Consistency of isotropic modified Maxwell theory: Microcausality and unitarity," Nucl. Phys. B **848**, 90 (2011), arXiv:1011.4258.

[5] F.R. Klinkhamer and M. Schreck, "Models for low-energy Lorentz violation in the photon sector: Addendum to 'Consistency of isotropic modified Maxwell theory'," Nucl. Phys. B **856**, 666 (2012), arXiv:1110.4101.

[6] S.R. Coleman and S.L. Glashow, "High-energy tests of Lorentz invariance," Phys. Rev. D **59**, 116008 (1999), arxiv:hep-ph/9812418.

[7] C. Kaufhold and F. R. Klinkhamer, "Vacuum Cherenkov radiation and photon triple-splitting in a Lorentz-noninvariant extension of quantum electrodynamics," Nucl. Phys. B **734**, 1 (2006), arXiv:hep-th/0508074.

[8] J.S. Diaz, F.R. Klinkhamer, and M. Risse, "Changes in extensive air showers from isotropic Lorentz violation in the photon sector," Phys. Rev. D **94**, 085025 (2016) , arXiv:1607.02099.

[9] B. de Wit and J. Smith, *Field Theory in Particle Physics, Volume 1* (North-Holland Publ., Amsterdam, 1986).